

2024-05-07
2023-0924

(Booklet)

Group	4	Herbicide
-------	---	-----------

IPCO®

CLOVITOX PLUS®

LIQUID HERBICIDE

CONTROLS MANY BROADLEAF WEEDS IN SEEDLING CLOVER, PEAS, PASTURE,
CEREALS, CORN AND SEEDLING GRASSES.

COMMERCIAL (AGRICULTURAL)

**READ THE LABEL AND BOOKLET BEFORE USING KEEP OUT OF
REACH OF CHILDREN**

ACTIVE INGREDIENTS: MCPB 375 g/L,
MCPA 25 g/L
(Both present as sodium salts)

**REGISTRATION NO. 24336
PEST CONTROL PRODUCTS ACT**

NET CONTENTS
10 litres

DANGER

[Insert octagon hazard symbol with skull and crossbones]
POISON

HARMFUL OR FATAL IF SWALLOWED; HARMFUL IF INHALED CORROSIVE TO
EYES

Interprovincial Cooperative Limited
P.O. Box 1050, Saskatoon, Saskatchewan S7K 3M9
1-204-233-3461

®T.M. Reg'd./M.D.

LABEL INDEX

Seedling Clover (Alsike, Ladino, Red, White Dutch, Wild White)	Section 1
Barley, Oats, Rye & Wheat	Section 2
Pasture	Section 3
Peas	Section 4
Field Corn	Section 5
Seedling Grasses	Section 6
Seedling Alfalfa/Seed Production	Section 7
Weeds Controlled / Rates of Application	Section 8
Mixing & Application Instructions	Section 9
Precautions	Section 10
First Aid	Section 11
Toxicological Information	Section 12
Rinsing and Disposal	Section 13

DIRECTIONS FOR USE:

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

SECTION 1

Crop: **SEEDLING CLOVER** (Alsike, Ladino, Red, White Dutch, Wild White) with or without a companion crop.

Note: Do not apply in hot weather (27°C or more) or under drought conditions.

Treatment Stage: Do not apply by air. Treat after primary or spade leaf to 4th true leaf stage. If a companion crop is seeded, make sure it is also at the correct stage of growth for safe treatment. See Barley, Oats, Rye and Wheat in Section 2.

Do not apply more than once per year.

GO TO SECTION 8

SECTION 2

Crop: **BARLEY, OATS, RYE, WHEAT**

Note: Do not apply in hot weather (27°C or more) or under drought conditions.

Treatment Stage: For ground or aerial application. Apply from 2 leaf to flag leaf stage.

Do not apply more than once per year.

GO TO SECTION 8

SECTION 3

Crop: **PASTURE**

Note: Do not apply in hot weather (27°C or more) or under drought conditions.

Treatment Stage: For ground or aerial application. Apply after grazing or cutting when weeds are at a susceptible stage and have sufficient leaf area to be contacted by spray.

Do not apply more than once per season.

GO TO SECTION 8

SECTION 4

Crop: **PEAS - PROCESSING OR FIELD TYPES**

Note: Do not apply in hot weather (27°C or more) or under drought conditions or when peas are under stress due to disease or other adverse field conditions. Do not apply more than once per year.

Treatment Stage: Do not apply by air. Apply when peas have 3-6 expanded leaves. Do not apply after the 6 leaf stage as damage may occur.

GO TO SECTION 8

SECTION 5

Crop: **FIELD CORN**

Note: Do not apply in hot weather (27°C or more) or under drought conditions.

Treatment Stage: Do not apply by air. Apply when crop is 45 cm high but before the beginning of tasseling. Use drop nozzles to spray weeds and avoid spraying corn leaves.

Do not apply more than once per year.

For hand harvesting of field corn, re-entry is not permitted until 15 days after application. As such, a pre-harvest interval (PHI) of 15 days after application is required.

GO TO SECTION 8

SECTION 6

Crop: **SEEDLING GRASSES:** Seedling smooth brome grass, meadow brome grass, creeping red fescue, reed canary grass, alai fescue, meadow fescue, Tall fescue, alai wild ryegrass, Russian wild ryegrass, Timothy, crested wheatgrass, intermediate wheatgrass, pubescent wheatgrass, tall wheatgrass, slender wheatgrass, streambank wheatgrass, northern wheatgrass, western wheatgrass, and green needlegrass.

Treatment Stage: Do not apply by air. Apply at the 2-4 leaf stage of the seedling grasses when weeds and crop are actively growing.

Do not apply more than once per season.

GO TO SECTION 8

SECTION 7

Crop: SEEDLING ALFALFA Grown for Seed Production (Western Canada Only)

NOTE TO USER: READ THE FOLLOWING BEFORE USING THIS PRODUCT FOR THE INDICATED SPECIAL USE APPLICATIONS.

The DIRECTIONS FOR USE for the uses described in this section of the label were developed by persons other than Interprovincial Cooperative Limited under the User Requested Minor Use Label Expansion program. For these uses, Interprovincial Cooperative Limited has not fully assessed performance (efficacy) and/or crop tolerance (phytotoxicity) under all environmental conditions or for all crop varieties when used in accordance with the label. The user should test the product on a small area first, under local conditions and using standard practices, to confirm the product is suitable for widespread application.

Treatment Stage: Do not apply by air. Apply at the 3 to 6 trifoliate leaf stage of seedling alfalfa when the Canada Thistle and other broadleaf weeds are at a susceptible stage and the crop is growing well. Alfalfa vigour may be reduced in the year of treatment, however, the crop recovers and yield will not normally be affected.

Do not apply more than once per season

GO TO SECTION 8

SECTION 8

WEEDS CONTROLLED/RATES OF APPLICATION

Apply when weeds are in the stage of growth indicated. Tolerance increases with age of crop and weeds.

A) WEEDS CONTROLLED

Annual Broadleaf Weeds:	Spray Stage:	Rate:
Wild Mustard, Ball Mustard, Worm Seed Mustard, Stinkweed, Lambs-quarter's	Spray in seedling stage.	2.75 L/ha
Redroot Pigweed, Ragweed, Shepherd's Purse, Volunteer Rapeseed (including Canola).	Spray in seedling stage.	4.25 L/ha

B) WEEDS SUPPRESSED

(i) Annual Broadleaf Weeds*:	Spray Stage:	Rate:
Wild Radish, Hempnettle, Annual Sow Thistle	Spray in seedling stage.	4.25 L/ha*

*NOTE: Control of this group of Annual Broadleaf Weeds may not be complete when treated under stress conditions such as hot weather, drought or if treated at more advanced stages.

(ii) Perennial Weeds*:	Spray Stage:	Rate:
Bull Thistle	Spray at rosette to early bud stage.	4.25 L/ha
Canada Thistle	Spray from 15 cm to early bud stage.	4.25 L/ha
Curled Dock	Spray at rosette stage.	4.25 L/ha
Plantains	Spray at rosette stage.	4.25 L/ha
Perennial Sow Thistle**	Spray in rosette stage.	4.25 L/ha
Field Bindweed**	Spray in spring.	4.25 L/ha
Horsetail	Spray when 15 cm high.	4.25 L/ha
Creeping Buttercup, Tall Buttercup	Spray in spring during rapid growth.	4.25 L/ha

*NOTE: Top growth control of Perennial Weeds only. Weeds will regrow by next growing season.

**NOTE: Application will retard flowering and growth of perennial sow thistle and check top growth of field bindweed.

Caution: Do not exceed 3.5 litres per hectare of product on seedling grasses.

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact Interprovincial Cooperative Ltd. at 1-204-233-3461 for information before applying any tank mix that is not specifically recommended on this label.

GO TO SECTION 9

SECTION 9

MIXING AND APPLICATION INSTRUCTIONS:

1. Fill spray tank 1/2 full with water. Start agitation.
2. Add the herbicide.
3. Complete filling spray tank with water.
4. Use 175-225L of water per ha for application by ground equipment.
5. **DO NOT USE LESS THAN 175 L OF WATER PER HA FOR GROUND APPLICATION.**
6. Do not exceed 275 kPa pressure for ground application

Field Sprayer Application: DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) coarse classification. Boom height must be 60 cm or less above the crop or ground.

Groundboom

For Mixers/Loaders/Applicators:

If handling less than 165 L of product in one day, wear a long-sleeved shirt, long pants, shoes plus socks, and chemical resistant gloves. Chemical resistant gloves are not required to be worn during groundboom application but are required for clean-up and repair. According to these provisions, do not handle more than 165 L of product in one day (for example, 39 ha at the maximum rate of 4.25 L /ha). These restrictions are in place to minimize exposure to individual workers. Application may need to be performed over multiple days or using multiple workers.

If handling more than 165 L product in one day, use a closed-system for mixing/loading while wearing coveralls over a long-sleeved shirt, long pants, shoes plus socks, and chemical resistant gloves. Also, use a closed-cab that provides both a physical barrier and respiratory protection (i.e. dust/mist filtering and/or vapour/gas purification system). The closed cab must have a chemical resistant barrier that totally surrounds the occupant and prevents contact with pesticides outside the cab. Chemical-resistant gloves are not required to be worn inside the closed cab, but have them ready for leaving the cab during calibration, repair or cleaning of equipment. According to these provisions, do not handle more than 720 L of product in one day (for example, 170 ha at the maximum rate of 4.25 L/ha). These restrictions are in place to minimize exposure to individual workers. Application may need to be performed over multiple days or using multiple workers.

Spot Treatment

For Mixers/Loaders/Applicators – Using manually-pressurized handheld equipment:

Wear coveralls over a long-sleeved shirt, long pants, shoes plus socks, and chemical resistant gloves. Do not handle more than 4 L of product in one day (for example, 0.9 ha at the maximum rate of 4.25 L/ha). These restrictions are in place to minimize exposure to individual workers. Application may need to be performed over multiple days or using multiple workers.

Aerial Application: DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply when wind speed is greater than 16 km/h at flying height at the site of application. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) coarse classification. Reduce drift caused by turbulent wingtip vortices. Nozzle distribution along the spray boom length MUST NOT exceed 65% of the wing- or rotorspan.

Mixers/Loaders:

Use a closed-system for mixing/loading while wearing coveralls over a long-sleeved shirt, long pants, shoes plus socks, and chemical resistant gloves.

Applicators:

Wear a long-sleeved shirt, long pants, and shoes plus socks.

Spray Buffer Zones

A spray buffer zone is NOT required for:
uses with hand-held application equipment permitted on this label.

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands), sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands), and estuarine/marine habitats.

Method of Application	Crop		Spray Buffer Zones (metres) Required for the Protection of:		
			Aquatic Habitat of Depths:		Terrestrial Habitat
			Less than 1 m	Greater than 1 m	
Field Sprayer	Peas, barley, field corn, oats, rye, seedling clover, wheat (spring and durum), seedling grasses, pasture and seedling alfalfa for seed		1	1	4
Aerial	Barley, oats, rye, wheat (spring and durum)	Fixed wing	1	0	175
		Rotary wing	1	0	125
Aerial	Pasture	Fixed wing	5	1	175
		Rotary wing	4	1	125

When tank mixes are permitted, consult the labels of the tank-mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

The spray buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Pesticides portion of the Canada.ca website

Use Precautions for Aerial Application (to cereal crops, pastures):

Apply only by fixed-wing or rotary aircraft equipment, which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label.

Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate recommended for aerial application on this label. Where no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment.

Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking devices

Use Precautions

Apply only when meteorological conditions at the treatment site allow for complete and even crop coverage. Apply only under conditions of good practice specific to aerial application as outlined in the National Aerial Pesticide Application Manual, developed by the Federal/Provincial/Territorial Committee on Pest Management and Pesticides.

Operator Precautions

Do not use human flaggers.

Do not allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of premixed chemicals with a closed system is permitted.

It is desirable that the pilot have communication capabilities at each treatment site at the time of application.

The field crew and the mixer/loaders must wear chemical resistant gloves, a long-sleeved shirt, long pants, coveralls, shoes plus socks, and goggles or face shield during mixing/loading, cleanup and repair. Applicators must wear long-sleeved shirt, long pants, and shoes plus socks. Follow the more stringent label precautions in cases where the operator precautions exceed the generic label recommendations on the existing ground boom label.

All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

Product Specific Precautions

Read and understand the entire label before opening this product. If you have questions, call the manufacturer at 1-204-233-3461 or obtain technical advice from the distributor or your provincial agricultural representative.

Application of this specific product must meet and/or conform to the following:

Volume: A minimum volume of 30 L per hectare of spray solution should be used. Use boom pressure of 235 kPa or less. Avoid placing nozzles where spray will enter wing tip vortices.

SECTION 10

PRECAUTIONS

Harmful or fatal if swallowed.

Harmful if inhaled. Avoid inhaling/breathing dusts, sprays, or other aerosolized particles.

1. **KEEP OUT OF REACH OF CHILDREN.** Do not use in residential areas, which are defined as sites where bystanders including children may be present during or after spraying, including around homes, schools, parks, playgrounds, playing fields, public buildings or any other areas where the general public including children could be exposed. Apply only to agricultural crops when the potential for drift to areas of human habitation and areas of human activity (houses, cottages, schools and recreational areas) is minimal. Take into consideration wind speed, wind direction, temperature inversion, application equipment and sprayer settings.
2. Wear a long-sleeved shirt, long pants, socks, shoes, goggles or face shield, suitable respirator and chemical-resistant gloves during mixing, loading, application and clean-up. Aerial applicators and applicators using a closed cab are not required to wear chemical-resistant gloves.
3. **WARNING POISON:** Harmful if absorbed through skin. **WARNING CORROSIVE:** Corrosive to eyes. Do not get in eyes, on skin or clothing. When handling, do not eat, drink or smoke. Take a shower daily, immediately after completing application. Wear freshly laundered clothes daily. Handle with care and mix only in a closed container. Use only in a well ventilated area.
4. Avoid spray or vapour drift onto susceptible plants such as tomatoes, tobacco, sugar beets, sunflowers, beans, turnips, cauliflower, cabbage, alfalfa, potatoes, ornamental trees, shrubs or flowers, canola, flax, or other desirable vegetation.
5. Do not spray when there is any danger of wind or on exceptionally hot days over 27°C.
6. Do not use in a greenhouse.
7. **Restricted Entry Intervals (REI):**
 - a. Cereals, pastures, seedling alfalfa, seedling clover, seedling grasses, peas (succulent/processing and dry/field): An REI of 12 hours after application is required to perform postapplication activities in treated areas.
 - b. Field corn: An REI of 9 days after application is required to perform postapplication activities (scouting) in treated areas.

A minimum rotational crop plant back interval (PBI) of 12 months must be observed for all crops other than those registered for use with MCPA or MCPB.

ENVIRONMENTAL PRECAUTIONS:

Toxic to aquatic organisms, birds, small wild animals, and non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.

Surface runoff

To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

Leaching

This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

STORAGE

Store this product away from food and feed. Keep in original container during storage.

RESISTANCE MANAGEMENT RECOMMENDATIONS:

For resistance management, IPCO Clovitox Plus Liquid Herbicide is a Group 4 herbicide. Any weed population may contain or develop plants naturally resistant to IPCO Clovitox Plus Liquid Herbicide and other Group 4 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

Where possible, rotate the use of IPCO CLOVITOX PLUS LIQUID HERBICIDE or other Group 4 herbicides within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.

Use tank mixtures with herbicides from a different group when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.

Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.

Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.

Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.

Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.

For further information or to report suspected resistance, contact Interprovincial Cooperative, Ltd. at 1-204-233-3461.

PRE-HARVEST INTERVALS:

Seedling forage grasses: Do not graze the treated crop or cut for hay in the year of establishment.

All other crops: Do not graze or cut for forage until 30 days after application. Withdraw meat animals from treated fields at least 3 days before slaughter.

GO TO SECTION 11

SECTION 11

FIRST AID:

1. If on skin or clothing, take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.
2. If in eyes, hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.
3. If swallowed, call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.
4. If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.
5. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

GO TO SECTION 12

SECTION 12

TOXICOLOGICAL INFORMATION:

High concentrations of MCPA or MCPB may cause severe irritation to the eyes. Symptoms of overexposure to MCPA or MCPB could include slurred speech, twitching, jerking and spasms, drooling, low blood pressure and unconsciousness. No specific antidote. Treat symptomatically. Symptomatic measures after ingestion include gastric lavage. Eye, skin and mucous membranes irritated due to alkaline nature of product.

GO TO SECTION 13

SECTION 13

DISPOSAL:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial/territorial requirements. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial/territorial regulatory agency. Contact the manufacturer and the provincial/territorial regulatory agency in case of a spill, and for clean up of spills.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

(Base Label)

Group	4	Herbicide
-------	---	-----------

IPCO®

CLOVITOX PLUS®

LIQUID HERBICIDE

CONTROLS MANY BROADLEAF WEEDS IN SEEDLING CLOVER, PEAS, PASTURE,
CEREALS, CORN AND SEEDLING GRASSES.

COMMERCIAL (AGRICULTURAL)

**READ THE LABEL AND ATTACHED BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN**

ACTIVE INGREDIENTS: MCPB 375 g /L,
MCPA 25 g /L
(Both present as sodium salts)

**REGISTRATION NO. 24336
PEST CONTROL PRODUCTS ACT**

**NET CONTENTS
10 litres**

DANGER
[Insert octagon hazard symbol with skull and crossbones]
POISON

WARNING POISON: HARMFUL OR FATAL IF SWALLOWED; HARMFUL IF INHALED
CORROSIVE TO EYES

Interprovincial Cooperative Limited
P.O. Box 1050, Saskatoon, Saskatchewan S7K 3M9
1-204-233-3461

®T.M. Reg'd./M.D.

(Back Panel)

IPCO CLOVITOX PLUS Liquid Herbicide

PRECAUTIONS:

Harmful or fatal if swallowed

Harmful if inhaled. Avoid inhaling/breathing dusts, sprays, or other aerosolized particles.

1. KEEP OUT OF REACH OF CHILDREN. Do not use in residential areas, which are defined as sites where bystanders including children may be present during or after spraying, including around homes, schools, parks, playgrounds, playing fields, public buildings or any other areas where the general public including children could be exposed. Apply only when the potential for drift to areas of human habitation or areas of human activity (houses, cottages, schools and recreational areas) is minimal. Take into consideration wind speed, wind direction, temperature inversion, application equipment and sprayer settings.
2. Wear a long-sleeved shirt, long pants, socks, shoes, goggles or face shield, suitable respirator and chemical-resistant gloves during mixing, loading, application and clean-up. Aerial applicators and applicators using a closed cab are not required to wear chemical-resistant gloves.
3. WARNING POISON: Harmful if absorbed through skin. WARNING CORROSIVE: Corrosive to eyes. Do not get in eyes, on skin or clothing. When handling, do not eat, drink or smoke. Take a shower daily, immediately after completing application. Wear freshly laundered clothes daily. Handle with care and mix only in a closed container. Use only in a well ventilated area.
4. Avoid spray or vapour drift onto susceptible plants such as tomatoes, tobacco, sugar beets, sunflowers, beans, turnips, cauliflower, cabbage, alfalfa, potatoes, ornamental trees, shrubs or flowers, canola, flax, or other desirable vegetation.
5. Do not use in a greenhouse.
6. .
7. Restricted Entry Intervals (REI):
 - c. Cereals, pastures, seedling alfalfa, seedling clover, seedling grasses, peas (succulent/processing and dry/field): An REI of 12 hours after application is required to perform postapplication activities in treated areas.
 - d. Field corn: An REI of 9 days after application is required to perform postapplication activities (scouting) in treated areas.

ENVIRONMENTAL PRECAUTIONS: Toxic to aquatic organisms, birds, small wild animals, and non-target terrestrial plants. Refer to the booklet for complete environmental precautions.

FIRST AID:

1. If on skin or clothing, take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.
2. If in eyes, hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.
3. If swallowed, call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.
4. If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.
5. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

High concentrations of MCPA or MCPB may cause severe irritation to the eyes. Symptoms of overexposure to MCPA or MCPB could include slurred speech, twitching, jerking and spasms, drooling, low blood pressure and unconsciousness. No specific antidote. Treat symptomatically. Symptomatic measures after ingestion include gastric lavage. Eye, skin and mucous membranes irritated due to alkaline nature of product.

STORAGE

Store this product away from food and feed. Keep in original container during storage.

DISPOSAL:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container

collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial/territorial requirements. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial/territorial regulatory agency. Contact the manufacturer and the provincial/territorial regulatory agency in case of a spill, and for clean up of spills.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.